Appln. No. 09/807,610 Amd. dated March 1, 2005 Reply to Office Action of September 2, 2004

Amendments to the Claims

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

1(Currently amended). An expression vector for the production of [[an]] a mixture of icIL-1ra-II protein proteins beginning at amino acid residue position +1 and +2 from the deduced start of translation on the icIL-1ra-II coding sequence, comprising a genomic DNA sequence encoding a human growth hormone signal peptide genomic DNA sequence, joined to a DNA segment encoding intracellular IL-1 receptor antagonist type II (icIL-1ra-II) and operably linked to a promoter sequence, whereby said icIL-1ra-II [[is]] proteins are expressed from said promoter sequence to produce [[an]] a mixture of icIL-1ra-II proteins with the amino acid sequence of SEQ ID NO:4 or SEQ ID NO:11 at the N-terminus.

Claim 2 (Cancelled).

3 (Previously presented). An isolated host cell line transformed with the expression vector of claim 1.

Claim 4 (Cancelled).

Appln. No. 09/807,610 Amd. dated March 1, 2005 Reply to Office Action of September 2, 2004

5 (Previously presented). An isolated host cell line in accordance with claim 3, wherein said cell is an endogenous cell of a human host.

Claim 6 (Cancelled).

7 (Previously presented). A method for producing a recombinant icIL-1ra-II comprising the steps of:

culturing a host cell line according to claim 3 to express and produce a recombinant glycosylated icIL-lra-II;

recovering the produced recombinant glycosylated icIL-1ra-II.

Claim 8 (Cancelled).

- 9 (Currently amended). An isolated glycosylated icIL1ra-II beginning at amino acid residue position +2 from the
 deduced start of translation on the icIL-1ra-II coding sequence
 and having the amino acid sequence of SEQ ID NO:11 at the Nterminus.
- 10 (Currently amended). The glycosylated icIL-1ra-II according to claim 9 having an apparent molecular weight of about 27 kDa on SDS-PAGE under reducing conditions with 15% acrylamide.

Appln. No. 09/807,610 Amd. dated March 1, 2005 Reply to Office Action of September 2, 2004

- 11 (Currently amended). The glycosylated icIL-1ra-II according to claim 9 having an apparent molecular weight of about 30 kDa on SDS-PAGE under reducing conditions with 15% acrylamide.
- 12 (Currently amended). A pharmaceutical composition, comprising the glycosylated icIL-1ra-II according to claim 9 in a therapeutically effective amount and a pharmaceutically acceptable excipient.
- 13 (Withdrawn). A method for reducing the amount of IL-1 in a patient having a condition associated with overexpression of IL-1, comprising administering the pharmaceutical composition according to claim 12 to a patient in need thereof.
- 14 (Withdrawn). A method for reducing the amount of IL-1 at a desired site in a human patient, comprising introducing a vector in accordance with claim 3 into appropriate endogenous human cells at the desired site to produce transformed cells which will express icIL-1ra-II at the desired site.

Claim 15-18 (Cancelled).